



Radiarel™

Benzylamine Mouthwash B. P. 0.15% w/v.

ALCOHOL FREE ANTI INFLAMMATORY ANALGESIC



RELIC
Biotechnology Pvt. Ltd.

An ISO 9001:2008 Certified Company

Patient Information

ABOUT your medicine

Benzydamine has analgesic, anti-inflammatory, antipyretic and anti-microbial properties. It is widely used in treating painful inflammation of mouth and throat conditions including tonsillitis, sore throat and mucositis.

HOW should I take the medicine?

For adults, rinse the mouth with 15ml of the solution 4 times a day at 6 hours interval or as recommended by the oncologist.

Gargle with the solution, or rinse around the mouth and throat, for at least 30 seconds, then spit out.

What should I do if I FORGET a dose?

Take the missed dose as soon as you remember. However, if it is almost time for your next dose, skip the dose you missed and take only your next scheduled dose. Do not double your dose to make up for a missed dose.

What PRECAUTIONS should I take?

This solution should not be given to children below the age of 12 years.

Inform your doctor if:

- You are allergic to any drugs.
- You have any other medical problems.

Be careful not to splash the solution into your eyes. If this happens, wash your eyes immediately with cold water.

Do not swallow the solution. If a normal 15ml dose is accidentally swallowed, you need not worry as it is unlikely to cause you any problem. If a large amount is swallowed, you should contact your doctor or go to the nearest hospital.

How should I STORE the medicine?

Store the medicine in a cool, dry place. Protect from moisture, heat and direct light.

Keep out of reach of children.

Once you have opened the mouthwash, do not keep it for more than 6 months.

Do not use if you notice any visible signs of deterioration of your medicine such as discoloration or the solution becoming cloudy. Return it to your pharmacist.

What are the possible SIDE-EFFECTS of the medicine?

Following the use as a mouth or throat preparation, oral numbness may happen. This is harmless and will soon disappear.

Occasionally, some people may feel a stinging sensation when they first use it. If this happens to you, try diluting your next dose with water to reduce the stinging.

Change in sense of taste, tingling, dryness or thirst and warm feeling of the mouth.